

# Perinatal Bundle- Elective Induction Bundle

## Data Collection Tool

**Elements:**

- Gestational Age 39 weeks or >.** Documented prior to initiation of oxytocin. Per ACOG definition in ACOG Practice Bulletin Number 10, 1999 {Induction of Labor}.

**Team Definition** \_\_\_\_\_

- Reassuring Fetal Status:** Assessed and documented prior to initiation of oxytocin *and* during administration.

**Team Definition** \_\_\_\_\_

- Pelvic Examination:** This element includes documentation of a complete pelvic assessment with cervical examination (dilation, effacement, station of the presenting part, cervical position and consistency; Bishop’s Score), clinical pelvimetry and an assessment of the fetal presentation.

**Team Definition** \_\_\_\_\_

- Hyperstimulation:** Absent at initiation of oxytocin and continuously assessed throughout augmentation. If present, it is recognized and treated.

**Team Definition** \_\_\_\_\_

**Team Definition** \_\_\_\_\_

**Instructions:** Review 5 charts each week where oxytocin was used to electively induce labor.

N: Total number of completed charts (pass- all elements must be present).

D: Total number of charts reviewed (5).

Month \_\_\_\_\_ Week \_\_\_\_\_

Chart	Gestational Age	Reassuring Fetal Status	Pelvic Examination	Hyperstimulation	Pass/Fail
#1					
#2					
#3					
#4					
#5					

Results= (N)Total Charts that Pass/ (D)Total Charts Reviewed.

