

HIIN – Venous Thromboembolism

New Jersey Hospital Association, Princeton

WEBINAR

EDU 2034W
Feb. 27, 2020
1 p.m. – 2 p.m.

REGISTRATION LINK:

<https://njha.webex.com/njha/onstage/g.php?MTID=e708f698b96743c2878395552f8d39c0b>

OVERVIEW:

Join us for a webinar on the latest evidence-based information for the prevention and management of Venous Thromboembolism (VTE) presented by Dr. Gregory A. Maynard.

Being in the hospital is a major risk factor for the development of venous thromboembolism (VTE). Patients with decreased mobility due to bedrest or recovery, or who experience blood vessel trauma due to surgery or other serious injury, are more likely to develop blood clots.

In fact, up to 60 percent of all VTE cases occur during or within 90 days of hospitalization, making it a leading preventable cause of hospital death.

To prevent VTE, hospital staff should evaluate patients for their risk of developing blood clots and use proper prevention and treatment procedures.

FACTS AND FIGURES:

- Up to 60 percent of VTE cases occur during or after hospitalization, making it a leading preventable cause of hospital death.
- In the U.K., upwards of 32,000 cases of hospital-associated VTE occur every year.
- In the U.S., more than 540,000 hospitalized patients develop VTE.
- In Australia, 30,000 cases of hospital-associated VTE occur every year.

TARGET AUDIENCE:

Physicians, nurse practitioners, registered nurses, pharmacists, PharmDs.

FACULTY:

Gregory A. Maynard, MD, MS, MHM
Chief Quality Officer
UC Davis Medical Center